

Ocular delivery

Arto Merivaara




INTEGRATING
Delivery Science
ACROSS DISCIPLINES



Absolute quantitation of barcoded liposomes in ocular tissues

11.7.2024


Arto Merivaara, PharmD
Postdoctoral researcher
Ocular Drug Delivery group
arto.merivaara@uef.fi







Barcoded drug delivery systems?

- Drug delivery systems (DDS) labeled with different tags that do not alter their properties
- Barcoding enables screening of multiple DDS within the same subject
- Rule of 3R's!
- Reducing the number of animals required
- Refinining the results





Why we need barcoding?





How can we do barcoding?

1. Barcode



DNA



RNA




Lipid



Peptide




2. Reader



LC-MS



PCR




Flow cytometry



Barcoding of liposomes

- DNA barcodes for a theranostic application: PCR analysis of DNA combined with live-dead cell imaging to study drug efficacy
- DNA/mRNA barcodes for LNP screening: Studying the efficacy of mRNA delivery for protein production






Absolute quantitation of barcoded liposomes in ocular tissues




AIM To develop a quantification method that allows studying of multiple liposome formulations within the same animal.

HYPOTHESIS Ocular pharmacokinetics of various liposome formulations can be studied within the same animal when the liposomes are labeled with barcode lipids.





Sample preparation and method validation




Merivaara, Arto, et al. *Journal of Controlled Release* 370 (2024): 1-13.




Concentration of barcode liposomes in ocular tissues

- Rats: intravitreal injections, injection volume 3 μ l
- 6 barcode liposome formulations in one injection
- Fluorescent label to confirm successful injection and to compare with barcode results
- Injection on day 0, tissues collected on day 0, day 1, day 4, and day 6.




A) Barcode liposomes in AH and Vitreous



Dose normalized concentrations



B) Barcode liposomes in neural retina



Merivaara, Arto, et al. *Journal of Controlled Release* 370 (2024): 1-13.



Evaluation of pharmacokinetics



Vitreous	50 nm PEG pos	50 nm PEG ionizable	100 nm PEG pos	100 nm PEG ionizable
Dose (ng)	74	106	74	162
C₀ (ng/ml)	1888	2571	1115	2822
K_e (day⁻¹)	0.211	0.277	0.160	0.187
T_{1/2} (day)	3.3	2.5	4.3	3.7
V_d (ml)	39	41	66	57
AUC(0-inf)	8959	9274	6982	15051
Aqueous humor	50 nm PEG pos	50 nm PEG ionizable	100 nm PEG pos	100 nm PEG ionizable
K_e (day⁻¹)	0.305	0.311	0.249	0.229
T_{1/2} (day)	2.3	2.2	2.8	3.0

Merivaara, Arto, et al. *Journal of Controlled Release* 370 (2024): 1-13.



Conclusions and take home message

- Barcode lipids offer an accurate, precise, and reliable method to study the in vivo intraocular pharmacokinetics of liposomes even in rats.
- The number of animals decreased from 72 to 12 by using the barcoded liposomes. This is important from an ethical point of view as well as from an economic point of view.
- The approach can be transferred to other tissues such as plasma and liver, to study the drug delivery with liposomes.






Acknowledgements

Prof. Arto Urtti's ODD group (UEF)

- Prof. Seppo Auriola, Prof. Tetsuya Terasaki, Dr. Kati-Sisko Vellonen, Dr. Tatu Lajunen, Dr. Jooseppi Puranen, Dr. Amir Sadeghi, Dr. Natalia Zashikhina, Lea Pirskanen, and all ODD group members!



Funders

- Academy of Finland Flagship project GeneCellNano
- Sigrid Juselius foundation
- Finnish cultural foundation
- Tandem Industry Academia foundation



Figures created with biorender.com.